

Correlation Between Sex Hormone Levels and Corneal Hysteresis and Resistance Factor During Menstrual Cycle

Mohammadreza Peyman, Azam Bakhtiari, Voon Pei Loo, Visvaraja Subrayan

The authors have no financial interests to disclose.


* Department of Ophthalmology, University Malaya, Kuala Lumpur, Malaysia

- * To determine corneal biomechanical properties change during phases of the menstrual cycle and to correlate with sex hormone levels..



* Purpose

* 24 eyes of 24 healthy young women (18-25 years old) were recruited. Hormones including follicle stimulating hormone (FSH; mU/ml), luteinizing hormone (LH; mU/ml), progesterone (ng/ml) and 17- β estradiol (pg/ml) were measured in serum three times in a cycle during the follicular (day 3 to day 9), luteal phases (day 21 to day 28) and ovulation (day 13 to day 15). At every time point, corneal hysteresis (CH) and the corneal resistance factor (CRF) were measured with the Ocular Response Analyzer.



Methods*

*The mean corneal hysteresis values at follicular, ovulatory, and luteal phases were 12.3mm Hg, 11.4mm Hg, and 12.7 mmHg ($p < 0.05$), and the mean corneal resistance factor values at the same time points were 10.3 9.7, and 10.6 mmHg ($p > 0.05$), respectively. There was a reverse correlation between serum estradiol and CH (Pearson correlation=-0.23, $P=0.041$)

Results*

*Biomechanical property of cornea changes during a menstrual cycle that might be due to serum estradiol levels.

Conclusion*

<http://www.drpeyman.net>

Department
Ophthalmology, University
Malaya ,
Kuala Lumpur, Malaysia

T: +60173387260

drmpeyman@yahoo.com

 **Contact**
Contact